



Product information



Lebosol[®]-Ammonium TS 200

Straight liquid inorganic macronutrient fertiliser

N (S) fertiliser solution 15 (+21)

15% Total nitrogen (200 g/l N) (9.3% Ammoniacal nitrogen; 5.7% Urea nitrogen)

21.3% Sulphur, water soluble (280 g/l S)

Crops with nutrient deficiency will be more susceptible against diseases and abiotic stress. Foliar fertilization with macro-and micro-elements will ensure an optimized plant nutrition.

Crop	Aim/Problem	Recommendation	Time
In all crops	For nitrogen and sulphur nutrition, vitality, leaf quality, yield, oil content, photosynthesis rate	5 l/ha ((in 200 – 400 l water) 0.5 – 1% for normal crop, 0.25% – 0.5% for sensitive crops under glass/film Do not use during flowering!)	When required
In all crops	For nitrogen and sulphur nutrition, vitality, leaf quality, yield, oil content, photosynthesis rate	Fertigation	Ask your consultant
Cereals	N efficiency, protein content, quality	2 – 4 times 5 l/ha	From 3-leaf-stage
Pasture land	N efficiency, vitality	2 – 4 times 5 l/ha	During the vegetation period
Potatoes	N efficiency, vitality	2 – 4 times 5 l/ha	From beginning of row closure
Legumes	N efficiency, increased vitality (e. g. in cold conditions), protein content	2 – 4 times 5 l/ha	From 6-leaf stage

Lebosol[®] Dünger GmbH

Wiesengasse 28 • 67471 Elmstein • Germany

Phone: +49 6328 98494-0

info@lebosol.de • www.lebosol.de/en

sustainable plant nutrition

Crop	Aim/Problem	Recommendation	Time
Maize	N efficiency, vitality	1 – 2 times 5 l/ha	From 4-leaf stage
Oilseed rape	N efficiency, vitality, oil content	2 – 4 times 5 l/ha	From 4-leaf stage
Sunflowers	N efficiency, vitality, oil content	1 – 2 times 5 l/ha	From 4-leaf stage
Sugar beet	N efficiency, vitality	2 – 4 times 5 l/ha	From 6-leaf stage
Medicinal plants, scented plants and spice plants	N efficiency, protein content, quality	2 – 4 times 5 l/ha	Once sufficient leaf mass has developed
General vegetables	N efficiency, protein content, quality	2 – 4 times 5 l/ha	Once sufficient leaf mass has developed
Hops	N efficiency, vitality	2 – 4 times 5 l/ha	From 0.5 m growth height
Greens	N efficiency, vitality	2 – 4 times 5 l/ha	During the vegetation period
Cotton	N efficiency, vitality	2 – 4 times 5 l/ha	From 4-leaf stage
Rice	N efficiency, protein content, quality	2 – 4 times 5 l/ha	From 3-leaf-stage

Lebosol® Dünger GmbH

Wiesengasse 28 • 67471 Elmstein • Germany

Phone: +49 6328 98494-0

info@lebosol.de • www.lebosol.de/en

sustainable plant nutrition 