



BIO

MADE IN GERMANY

Lebosol

Product information

Lebosol[®] - Boron

Straight inorganic micronutrient fertiliser

Micronutrient solution fertiliser

11% Boron as water soluble boron (150 g/l B)

also contains: 3.46% nitrogen (N)

Crops with nutrient deficiency will be more susceptible against diseases and abiotic stress.

Foliar fertilization with macro-and micro-elements will ensure an optimized plant nutrition.

Crop	Aim/Problem	Recommendation	Time
In all crops	For boron nutrition, calcium transport, pollen quality, yield, flowering quality, fruit quality, photosynthesis rate	0,5 – 3 l/ha (as foliar fertilization in 200 – 400 l water. Upon application with backpack sprayer 0.5%.)	When required
In all crops	Increased vitality (e. g. during frost for flowering)	1 – 2 times 1 l/ha (Best effect in combination with 2 l/ ha Aminosol [®] + 5 l/ha Lebosol [®] -Potassium 450)	When required
In all crops	For boron nutrition, calcium transport, pollen quality, yield, flowering quality, fruit quality, photosynthesis rate	Fertigation	Ask your consultant
Cereals	N efficiency, stem stability, seed quality, winter hardiness	1 – 2 times 0,5 – 1 l/ha	From 3-leaf-stage
Pasture land	N efficiency, vitality, tillering, stem stability, winter hardiness	1 – 4 times 0,5 – 1 l/ha	During the vegetation period

Lebosol[®] Dünger GmbH

Wiesengasse 28 • 67471 Elmstein • Germany

Phone: +49 6328 98494-0

info@lebosol.de • www.lebosol.de/en

sustainable plant nutrition



Crop	Aim/Problem	Recommendation	Time
Potatoes	N efficiency, internal quality, skin firmness	1 – 4 times 1 l/ha	From 6-leaf stage
Legumes	Yield, N efficiency, increased vitality (e. g. in cold conditions), protein content	2 times 1 l/ha	From 4-leaf stage
Maize	Energy density, N efficiency, yield, increased vitality (e. g. in cold conditions)	1 – 2 times 2 l/ha	From 4-leaf stage
Oilseed rape	Stem stability, husk firmness, N efficiency, winter hardiness, even maturation, oil content	2 – 4 times 2 – 3 l/ha	From 4-leaf stage
Sunflowers	Stem stability, N efficiency, increased vitality (e. g. in cold conditions), even maturation, oil content	1 – 2 times 2 l/ha	From 4-leaf stage
Sugar beet	N efficiency, increased vitality (e. g. in cold conditions), to prevent heart rot and dry rot, sugar content	2 – 3 times 2 – 3 l/ha	From 6-leaf stage
Strawberries	To prevent deformed berries; sugar content, flower bud development, winter hardiness	1 – 3 times 1 – 2 l/ha	From green buds
Strawberries	Plant quality in seed production crops: Strong plants, formation of offshoots	2 times 1 l/ha	14 and 7 days before grubbing up the young plants
Pome fruit	Fruit set, skin quality, flower bud development, winter hardiness	2 – 4 times 1 l/ha	From red buds
Stone fruit	Fruit set, skin quality, flower bud development, winter hardiness	2 – 3 times 1 l/ha	From fruit set
Soft fruit	Fruit set, skin quality, flower bud development, winter hardiness	1 – 3 times 1 l/ha	Start of shoot growth
Dessert grapes	Fruit set, skin quality, flower bud development, winter hardiness, even maturation	2 – 3 times 1 l/ha	From the enlargement of the inflorescences
Citrus fruits	Fruit set, skin quality, flower bud development, winter hardiness	2 – 4 times 1 l/ha	From white buds
Nuts	Fruit set, skin quality, flower bud development, winter hardiness	1 – 3 times 1 – 2 l/ha	From swelling buds

Lebosol® Dünger GmbH

Wiesengasse 28 • 67471 Elmstein • Germany

Phone: +49 6328 98494-0

info@lebosol.de • www.lebosol.de/en

sustainable plant nutrition 

Crop	Aim/Problem	Recommendation	Time
Wine grapes	Fruit set, skin quality, flower bud development, winter hardiness, even maturation	2 – 3 times 1 l/ha	From the enlargement of the inflorescences
General vegetables	Leaf quality, stem stability, N efficiency, vitality (e. g. in cold conditions), even maturation, oil content	2 – 4 times 1 – 3 l/ha	Once sufficient leaf mass has developed
Medicinal plants, scented plants and spice plants	Leaf quality, stem stability, N efficiency, vitality (e. g. in cold conditions), even maturation, oil content	2 – 4 times 1 – 3 l/ha	Once sufficient leaf mass has developed
Hops	Bud and shoot development, quality	3 – 5 times 1 – 2 l/ha	From 0.5 m growth height
Tobacco	N efficiency, leaf quality, vitality	1 – 3 times 1 – 2 l/ha	From 4-leaf stage
Christmas trees	Needle quality, N efficiency, vitality, winter hardiness	1 – 2 times 1 l/ha	From budding
Ornamental plants	N efficiency, leaf quality, vitality	1 – 2 times 1 l/ha	Once sufficient leaf mass has developed
Greens	N efficiency, vitality, stem stability, frost resistance	1 – 4 times 0,5 – 1 l/ha	During the vegetation period
Cotton	Fruit set, flower bud development, winter hardiness	1 – 3 times 1 – 2 l/ha	From budding
Rice	N efficiency, stem stability, seed quality, increased vitality (e. g. in cold conditions)	1 – 2 times 0,5 – 0,75 l/ha	From 3-leaf-stage

Lebosol® Dünger GmbH

Wiesengasse 28 • 67471 Elmstein • Germany

Phone: +49 6328 98494-0

info@lebosol.de • www.lebosol.de/en

sustainable plant nutrition 